## **CORRECTED AND SUBSTITUTE**

## AMENDMENT TO THE CLAIMS (filed October 13, 2004)

- 1-8 (Canceled)
- 9. (Currently Amended) A method of for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

exposing the damaged tissue, which is surrounded by an air impermeable wound cover, to an effective amount of gaseous nitric oxide for a period of time that exceeds eight consecutive hours;

allowing at least a portion of the gaseous nitric oxide to contact the air adjacent to the air impermeable would wound cover through the air impermeable wound cover; and

pretreating the damaged tissue with an agent in combination with gaseous nitric oxide in order to enhance its effectiveness and/or absorption.

10. (Currently Amended) A method of for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

exposing the damaged tissue, which is surrounded by an air impermeable wound cover, to an effective amount of gaseous nitric oxide for a period of time that exceeds eight consecutive hours;

allowing at least a portion of the gaseous nitric oxide to contact the air adjacent to the air impermeable would wound cover through the air impermeable wound cover; and

wetting, dampening, or moistening the damaged tissue following gaseous nitric oxide therapy.

11. (Currently Amended) A method of for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

exposing the damaged tissue, which is surrounded by an air impermeable wound cover, to an effective amount of gaseous nitric oxide for a period of time that exceeds eight consecutive hours;

allowing at least a portion of the gaseous nitric oxide to contact the air adjacent to the air impermeable would wound cover through the air impermeable wound cover; and

posttreating the damaged tissue by applying a wound healing agent in combination to gaseous nitric oxide therapy.

12. (Currently Amended) A method of for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

exposing the damaged tissue, which is surrounded by an air impermeable wound cover, to an effective amount of gaseous nitric oxide for a period of time that exceeds eight consecutive hours;

allowing at least a portion of the gaseous nitric oxide to contact the air adjacent to the air impermeable would wound cover through the air impermeable wound cover; and

posttreating the damaged tissue with an agent in combination with gaseous nitric oxide in order to enhance its effectiveness and/or absorption.

13. (Previously Presented) A method of for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

exposing the damaged tissue, which is surrounded by an air impermeable wound cover, to an effective amount of gaseous nitric oxide for a period of time that exceeds eight consecutive hours; and

administering exogenous nitric oxide to tissue flap and surrounding damaged area in order to promote flap viability and increase local blood flow to donated tissue.

14. (Original) A method for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

spraying, from a spray container, the damaged tissue with an effective amount of gaseous nitric oxide; and

allowing the gaseous nitric oxide to contact the air adjacent to the damaged tissue.

15. (Currently Amended) A method for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

exposing the damaged tissue, which is surrounded by an air impermeable wound cover, to an effective amount of gaseous nitric oxide; and

pretreating the damaged tissue with a wound healing agent other than gaseous nitric oxide.

- 16. (Canceled)
- 17. (Previously Presented) The method of claim 9 wherein the air impermeable wound cover is transparent and allows for permeation of small molecules, while simultaneously preventing microbial contamination of the damaged tissue from a source outside of the wound cover.
- 18. (Previously Presented) The method of claim 9 wherein the effective amount of gaseous nitric oxide ranges from 20-1000 ppm.
- 19. (Previously Presented) The method of claim 18 wherein the effective amount of gaseous nitric oxide is at least 200 ppm.
- 20. (Previously Presented) The method of claim 18 wherein the effective amount of gaseous nitric oxide is at least 400 ppm.
- 21. (Previously Presented) The method of claim 9 wherein the step of pretreating the damaged tissue comprises exposing the damaged tissue directly to the agent in combination with gaseous nitric oxide.

- 22. (Previously Presented) The method of claim 9 wherein the step of pretreating the damaged tissue comprises administering an agent to the patient that indirectly enhance the local amount of nitric oxide.
- 23. (Currently Amended) The method of claim <u>15</u> <del>16</del>, wherein the damaged tissue is selected from the group consisting of muscle, ligament, tendon, skin, bone, and cornea.
- 24. (Currently Amended) The method of claim <u>15</u> <del>16</del> wherein the damaged tissue is damaged by surgical incisions, trauma, and pathological processes.
- 25. (Currently Amended) The method of claim 15 16 wherein the effective amount of nitric oxide is at least 200 ppm.
- 26. (Currently Amended) The method of claim 15 16 wherein the effective amount of nitric oxide is at least 400 ppm.

## **CORRECTED AND SUBSTITUTE**

## **AMENDMENT TO THE CLAIMS (Filed September 21, 2004)**

- 1-8 (Canceled)
- 9. (Currently Amended) A method of for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

exposing the damaged tissue, which is surrounded by an air impermeable wound cover, to an effective amount of gaseous nitric oxide for a period of time that exceeds eight consecutive hours;

allowing at least a portion of the gaseous nitric oxide to contact the air adjacent to the air impermeable would cover through the air impermeable wound cover; and

The method of claim 1 further comprising pretreatement step of exposing pretreating the damaged tissue to with an agent in combination with gaseous nitric oxide in order to enhance its effectiveness and/or absorption.

10. (Currently Amended) A method of for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

exposing the damaged tissue, which is surrounded by an air impermeable wound cover, to an effective amount of gaseous nitric oxide for a period of time that exceeds eight consecutive hours;

allowing at least a portion of the gaseous nitric oxide to contact the air adjacent to the air impermeable would cover through the air impermeable wound cover; and

The method of claim 1 further comprising posttreatment step of wetting, dampening, or moistening the damaged tissue following gaseous nitric oxide therapy.

11. (Currently Amended) A method of for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

exposing the damaged tissue, which is surrounded by an air impermeable wound cover, to an effective amount of gaseous nitric oxide for a period of time that exceeds eight consecutive hours;

allowing at least a portion of the gaseous nitric oxide to contact the air adjacent to the air impermeable would cover through the air impermeable wound cover; and

The method of claim 1 further comprising postreatment step of posttreating the damaged tissue by applying a wound healing agent in combination to gaseous nitric oxide therapy.

12. (Currently Amended) A method of for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

exposing the damaged tissue, which is surrounded by an air impermeable wound cover, to an effective amount of gaseous nitric oxide for a period of time that exceeds eight consecutive hours;

allowing at least a portion of the gaseous nitric oxide to contact the air adjacent to the air impermeable would cover through the air impermeable wound cover; and

The method of claim 1 further comprising posttreatment step of exposing postreating the damaged tissue to with an agent in combination with gaseous nitric oxide in order to enhance its effectiveness and/or absorption.

13. (Currently Amended) A method of for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

exposing the damaged tissue, which is surrounded by an air impermeable wound cover, to an effective amount of gaseous nitric oxide for a period of time that exceeds eight consecutive hours; and

The method of claim 1 further comprising the administration of administering exogenous nitric oxide to tissue flap and surrounding damaged area in order to promote flap viability and increase local blood flow to donated tissue.

14. (Original) A method for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

spraying, from a spray container, the damaged tissue with an effective amount of gaseous nitric oxide; and

allowing the gaseous nitric oxide to contact the air adjacent to the damaged tissue.

15. (Currently Amended) A method for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

exposing the damaged tissue, which is surrounded by an air impermeable wound cover, to an effective amount of gaseous nitric oxide for a period of time that exceeds eight consecutive hours;

pretreating the damaged tissue with a wound healing agent other than gaseous nitric oxide.

16. (Original) A method for promoting the healing of damaged tissue in a patient in need of such treatment, comprising:

exposing the damaged tissue, which is surrounded by an air impermeable wound cover, to an effective amount of gaseous nitric oxide and other gases which contain oxygen for a period of time.

- 17. (New) The method of claim 9 wherein the air impermeable wound cover is transparent and allows for permeation of small molecules, while simultaneously preventing microbial contamination of the damaged tissue from a source outside of the wound cover.
- 18. (New) The method of claim 9 wherein the effective amount of gaseous nitric oxide ranges from 20-1000 ppm.
- 19. (New) The method of claim 18 wherein the effective amount of gaseous nitric oxide is at least 200 ppm.
- 20. (New) The method of claim 18 wherein the effective amount of gaseous nitric oxide is at least 400 ppm.

- 21. (New) The method of claim 9 wherein the step of pretreating the damaged tissue comprises exposing the damaged tissue directly to the agent in combination with gaseous nitric oxide.
- 22. (New) The method of claim 9 wherein the step of pretreating the damaged tissue comprises administering an agent to the patient that indirectly enhance the local amount of nitric oxide.
- 23. (New) The method of claim 16, wherein the damaged tissue is selected from the group consisting of muscle, ligament, tendon, skin, bone, and cornea.
- 24. (New) The method of claim 16 wherein the damaged tissue is damaged by surgical incisions, trauma, and pathological processes.
- 25. (New) The method of claim 16 wherein the effective amount of nitric oxide is at least 200 ppm.
- 26. (New) The method of claim 16 wherein the effective amount of nitric oxide is at least 400 ppm.